



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2020

---

## **The global burden of gastro-oesophageal reflux disease: more than just heartburn and regurgitation**

Sweis, Rami ; Fox, Mark

DOI: [https://doi.org/10.1016/S2468-1253\(20\)30002-9](https://doi.org/10.1016/S2468-1253(20)30002-9)

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-192837>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Sweis, Rami; Fox, Mark (2020). The global burden of gastro-oesophageal reflux disease: more than just heartburn and regurgitation. *Lancet Gastroenterology Hepatology*, 5(6):519-521.

DOI: [https://doi.org/10.1016/S2468-1253\(20\)30002-9](https://doi.org/10.1016/S2468-1253(20)30002-9)

## Achalasia: When a simple disease becomes complex

David A. Katzka MD, Mark Fox MD MA

### Author affiliations:

<sup>1</sup> Division of Gastroenterology and Hepatology, Mayo 9, Mayo Clinic, 200 First Ave., S.W. Rochester, MN 55905, USA

<sup>2</sup> Division of Gastroenterology and Hepatology, University Hospital Zürich, Zürich, Switzerland.

<sup>3</sup> Digestive Function: Basel, Laboratory and Clinic for motility disorders and functional Gastrointestinal diseases, Centre for integrative Gastroenterology, Klinik Arlesheim, Arlesheim, Switzerland.

### Corresponding author:

Prof. Dr. Mark Fox

Digestive Function: Basel, Laboratory and Clinic for Motility Disorders and Functional GI Diseases, Klinik Arlesheim, Arlesheim, Switzerland

Email: [dr.mark.fox@gmail.com](mailto:dr.mark.fox@gmail.com)

Tel: +41 79 1934795

## Editorial

Textbooks present achalasia as a relatively simple disease defined by esophageal aperistalsis and impaired “relaxation pressure” of the EGJ with radiographic findings that typically accompany these manometric findings. The etiology of this disorder is not known but cases have been linked to an autoimmune process limited to the myenteric plexus of the lower esophageal sphincter (LES) and initiated by molecular mimicry of a viral epitope, such as HSV.<sup>1,2</sup> Recent recognition that certain types of achalasia have esophageal spasm or even preserved peristalsis (EGJ outlet obstruction) has broadened the definition of the disease.<sup>3,4</sup> At the same time the wide spectrum of histopathologic findings - from complete neuronal loss to lymphocytic inflammation to apparently normal histopathology - emphasizes that “achalasia” is a heterogeneous condition.<sup>3,5</sup> There are also a plethora of secondary causes, including genetic conditions, paraneoplastic syndromes,<sup>6</sup> Chagas Disease<sup>7</sup> and opioid induced esophageal dysmotility.<sup>8</sup> Additionally, both benign inflammatory and malignant processes that involve the EGJ may produce findings similar to achalasia. Achalasia can be viewed as one disease with multiple potential etiologies, with the commonality being EGJ outlet obstruction.

The EGJ has a complex functional anatomy that includes an intrinsic physiological sphincter of smooth-muscle within the abdominal oesophagus [LES] and gastric cardia (clasp and sling fibers) together with an extrinsic crural diaphragm (CD) of skeletal muscle (**Figure HV panel**).<sup>9</sup> Previously the focus of attention in achalasia has been almost entirely on impaired LES function, although impaired relaxation of the proximal stomach has been documented.<sup>10,11</sup>

In this edition of *Gastroenterology*, researchers led by Ravi Mittal combined 3-dimensional high-resolution manometry (3D-HRM) with detailed analysis of computed tomography images to study the esophago-gastric junction (EGJ) in patients with achalasia and healthy controls.<sup>(Mittal reference)</sup> This novel contribution describes possible abnormalities of the esophageal hiatus and CD muscle in idiopathic achalasia.<sup>(Mittal reference)</sup> Consistent with previous reports,<sup>12</sup> the authors demonstrated an asymmetric, extrinsic CD component to EGJ pressure superimposed on the intrinsic LES. In all subjects, the LES length was greater on the lesser curvature as compared to the greater curvature of the stomach; whereas, LES related pressures were greater towards the greater curvature, at the esophago-gastric insertion angle - termed angle of His in anatomic studies (**Figure HV panel**). In achalasia patients the LES pressure distribution and the location of CD contraction within the EGJ high pressure zone were more variable and, in many cases, appeared to be abnormal. The authors attributed these changes to the asymmetric pull of the “noose-shaped” right crura on the abdominal esophagus.

Although innovative, the measurements derived from this methodology have not been validated and the soft-tissue resolution of CT to differentiate EGJ structures is at the limit of what is possible even with Magnetic Resonance (MR) Imaging.<sup>13</sup> Further, the mechanistic inferences are speculative and not necessarily consistent with published literature or clinical experience. Far from the LES being “tightly anchored” to the hiatus, striking shortening of the esophagus with pan-esophageal pressurization mediated by longitudinal muscle contraction is observed in achalasia patients and, when this occurs, the LES can be drawn into the chest while the diaphragmatic pinch remains in position!<sup>14,15</sup> Further,

surgeons do not report that the LES is neither firmly attached to the CD nor trapped within the hiatus in patients referred for Heller Myotomy (personal communication, Prof. Robert Rosenberg). However, repeated "pulling" on the phreno-esophageal ligaments during esophageal shortening could explain the physical breaks (muscle tears?) in the left crus of the diaphragm observed in the paper.

It is also clear that observational data cannot distinguish cause from effect. Choosing appropriate interventions to test the hypothesis is a challenge; however, Brasseur et al. have shown that it can be done by "pharmacological dissection".<sup>9</sup> Mittal et al. refer to a case in which injection of botulinum toxin into the sphincter region inadvertently paralyzed the CD and obliterated the esophago-gastric insertion angle. As botulinum toxin produces profound relaxation in both smooth and striated muscle, the effects of injection into the LES, gastric cardia or CD could produce the same result and, therefore, cannot be used to differentiate biophysical effects of these structures. Studies that combined MR imaging and HRM to assess EGJ structure and function in gastro-esophageal reflux disease (GERD) may provide more insight.<sup>16, 17</sup> The angle of His is more obtuse in GERD patients in healthy controls,<sup>17</sup> which is interesting to contrast with the acute insertion angle seen in achalasia patients ([Figure Achalasia and GERD panel](#)). This observation could be explained by a wide hiatus or weak CD function, both common in GERD patients; however, anatomical studies indicate that the clasp and sling fibers at the gastric cardia also maintain the angle of His. The biophysical mechanism of the EGJ reflux barrier was investigated by repeating measurements after treatment with baclofen. This GABA-B agonist not only restored the insertion angle in GERD patients but also reduced transient LES relaxations and reflux events.<sup>16</sup> Baclofen increases tone in the LES and the clasp and sling muscle fibers via effects on the vagal nerve;<sup>18</sup> however, it relaxes the crural diaphragm... an effect that explains its utility in hiccups!<sup>19</sup> Together, the results support the hypothesis that the acute insertion angle in achalasia and the obtuse insertion angle in GERD patients are related to variation in muscle tone within the clasp and sling fibers of the proximal stomach and not, or not only, with abnormal hiatal anatomy or CD function ([Figure](#)).

Another issue is the part of the hypothesis that suggests that degenerative changes in the spine can impact on CD function is unlikely to apply in younger patients with achalasia. Although previous work has demonstrated neurodegeneration secondary to obstruction in an animal model of achalasia, this etiology is hard to reconcile with evidence of autoimmune etiology in many patients.<sup>2, 5</sup>

Achalasia may once have appeared to be a simple motility disorder; however, it is now clear that many etiologies result in a common presentation that responds to treatment that relieves EGJ outlet obstruction. It is hoped that improved understanding of the disease will lead to individual treatment directed at the underlying pathophysiology. With researchers such as Mittal et al. prepared to develop innovative methods and present novel findings, this goal is getting closer.

## Figure Legend

Schematic diagram of esophago-gastric junction (EGJ) anatomy in health (central panel). Immediately below the lower esophageal sphincter, clasp and sling fibres fanning out from the greater to the lesser curvature form the distal component of the esophago-gastric junction high pressure zone. Increased density of these fibers on the greater than the lesser curvature could explain the variation in length and pressure of the sphincter between these two positions observed on 3D-HRM (lower panel). This anatomical arrangement may also maintain the esophago-gastric insertion angle. In achalasia patients with high tone in both components of the intrinsic sphincter the pressure is abnormally high and the insertion angle is acute (left panel). In GERD patients with low tone in the intrinsic sphincter the pressure is low and the angle is obtuse (right panel); however, administration of the GABA-B agonist baclofen increases pressure and restores the angle towards normal values. An alternative hypothesis proposed by Mittal et al. in this edition of *Gastroenterology* states that the changes in the structure and function of the crural diaphragm produces the effects seen on 3D-HRM and variation in the esophago-gastric insertion angle in health and disease.

## References:

1. Facco M, Brun P, Baesso I, et al. T cells in the myenteric plexus of achalasia patients show a skewed TCR repertoire and react to HSV-1 antigens. *Am J Gastroenterol* 2008;103:1598-609.
2. Clark SB, Rice TW, Tubbs RR, et al. The nature of the myenteric infiltrate in achalasia: an immunohistochemical analysis. *Am J Surg Pathol* 2000;24:1153-8.
3. Kahrilas PJ, Boeckxstaens G. The spectrum of achalasia: lessons from studies of pathophysiology and high-resolution manometry. *Gastroenterology* 2013;145:954-65.
4. Kahrilas PJ, Bredenoord AJ, Fox M, et al. Expert consensus document: Advances in the management of oesophageal motility disorders in the era of high-resolution manometry: a focus on achalasia syndromes. *Nat Rev Gastroenterol Hepatol* 2017;14:677-688.
5. Sodikoff JB, Lo AA, Shetuni BB, et al. Histopathologic patterns among achalasia subtypes. *Neurogastroenterol Motil* 2016;28:139-45.
6. Katzka DA, Farrugia G, Arora AS. Achalasia secondary to neoplasia: a disease with a changing differential diagnosis. *Dis Esophagus* 2012;25:331-6.
7. Goin JC, Sterin-Borda L, Bilder CR, et al. Functional implications of circulating muscarinic cholinergic receptor autoantibodies in chagasic patients with achalasia. *Gastroenterology* 1999;117:798-805.
8. Ratnapli SK, Crowell MD, DiBaise JK, et al. Opioid-Induced Esophageal Dysfunction (OIED) in Patients on Chronic Opioids. *Am J Gastroenterol* 2015;110:979-84.
9. Brasseur JG, Ulerich R, Dai Q, et al. Pharmacological dissection of the human gastro-oesophageal segment into three sphincteric components. *J Physiol* 2007;580:961-75.
10. Mearin F, Papo M, Malagelada JR. Impaired gastric relaxation in patients with achalasia. *Gut* 1995;36:363-8.
11. Bittinger M, Barnert J, Eberl T, et al. Postprandial gastric relaxation in achalasia. *Eur J Gastroenterol Hepatol* 1998;10:741-4.
12. Kwiatek MA, Pandolfino JE, Kahrilas PJ. 3D-high resolution manometry of the esophagogastric junction. *Neurogastroenterol Motil* 2011;23:e461-9.

13. Curcic J, Fox M, Kaufman E, et al. Gastroesophageal junction: structure and function as assessed by using MR imaging. *Radiology* 2010;257:115-24.
14. Hong SJ, Bhargava V, Jiang Y, et al. A unique esophageal motor pattern that involves longitudinal muscles is responsible for emptying in achalasia esophagus. *Gastroenterology* 2011;139:102-11.
15. Tutuian R, Pohl D, Castell DO, et al. Clearance mechanisms of the aperistaltic oesophagus: the "pump gun" hypothesis. *Gut* 2006;55:584-5.
16. Curcic J, Schwizer A, Kaufman E, et al. Effects of baclofen on the functional anatomy of the oesophago-gastric junction and proximal stomach in healthy volunteers and patients with GERD assessed by magnetic resonance imaging and high-resolution manometry: a randomised controlled double-blind study. *Aliment Pharmacol Ther* 2014;40:1230-40.
17. Curcic J, Roy S, Schwizer A, et al. Abnormal structure and function of the esophagogastric junction and proximal stomach in gastroesophageal reflux disease. *Am J Gastroenterol* 2014;109:658-67.
18. Smid SD, Young RL, Cooper NJ, et al. GABA(B)R expressed on vagal afferent neurones inhibit gastric mechanosensitivity in ferret proximal stomach. *Am J Physiol Gastrointest Liver Physiol* 2001;281:G1494-501.
19. Steger M, Schneemann M, Fox M. Systemic review: the pathogenesis and pharmacological treatment of hiccups. *Aliment Pharmacol Ther* 2015;42:1037-50.

